

# ECVP/ESVP Summer School in Veterinary Pathology



Marie Curie Training Courses


Summer School 2007 – Avian B2

### B2. Hen.

Organ? Morphologic diagnosis? Etiologic diagnosis?




890989



Organ? Keywords? Morphologic diagnosis? Etiologic diagnosis?

### B2. Hen




#### Liver

The architecture of the liver is markedly distorted by random, multifocal to coalescing, up to 1-mm diameter, variably encapsulated, granulomas.




890989



The larger granulomas contain, from the centre to the periphery: a central mass of amorphous hypereosinophilic material with karyorrhectic and pyknotic nuclear debris (caseous necrosis); cholesterol clefts;


## B2. Hen



... surrounded by a thick rim of large, pale acidophilic cells with indistinct borders and round hypochromatic nucleus (epithelioid macrophages); giant multinucleate cells with abundant eosinophilic cytoplasm and either peripheral nuclei (Langhans-type giant cells) or haphazardly-placed nuclei (foreign-body-type giant cells);




890989




... moderate amount of **lymphocytes**, few **plasma cells** and **heterophils**; a thin rim of **fibrosis**. Within granulomas, some macrophages are round, large, and filled with an empty cytoplasmic vacuole (**lipid-laden macrophages**).


### B2. Hen



Smaller granulomas contain **epithelioid macrophages**, Langhans giant cells, foreign body giant cells, and do not possess a necrotic centre.




890989



#### Fite-Farraco stain

Acid-fast bacilli within the necrotic centres of the granulomas, low amount






Within the remnant hepatic parenchyma, there are multiple foci of extramedullary heterophilic granulopoiesis (especially periportal) and lymphopoiesis ± amyloid deposition.



890989



Multifocal to coalescing marked granulomatous hepatitis and perihepatitis with caseous necrosis.

Mycobacterial hepatitis and perihepatitis.